MAP MF-1648-A

DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

MINERAL RESOURCE POTENTIAL MAP OF THE POND MOUNTAIN AND POND MOUNTAIN ADDITION ROADLESS AREAS, CARTER COUNTY, TENNESSEE

By W.R. Griffitts, M.L. Chatman, and R.F. Bitar

MISCELLANEOUS FIELD STUDIES MAP
Published by the U.S. Geological Survey, 1989

NEW YORK
JUL 5 1989
STATE LIBRARY

GRIFFITTS AND OTHERS—MINERAL RESOURCE POTENTIAL, POND MTN. AND POND MTN. ADDITION ROADLESS AREAS, TENNESSEE 1:24,000 MAP MF-1648-A

best copy available

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

MISCELLANEOUS FIELD STUDIES
MAP MF-1648-A
PAMPHLET

MINERAL RESOURCE POTENTIAL OF THE POND MOUNTAIN AND POND MOUNTAIN ADDITION ROADLESS AREAS, CARTER COUNTY, TENNESSEE

By

**W. R. Griffitts, U.S. Geological Survey,
and R. F. Bitar
and
M. L. Chatman, U.S. Bureau of Mines**

STUDIES RELATED TO WILDERNESS

Under the provisions of the Wilderness Act (Public Law 88-577, September 3, 1964) and related acts, the U.S. Geological Survey and the U.S. Bureau of Mines have been conducting mineral surveys of wilderness and primitive areas. Areas officially designated as "wilderness," "wild," or "canoe" when the act was passed were incorporated into the National Wilderness Preservation System, and some of them are presently being studied. The act provided that areas under consideration for wilderness designation should be studied for suitability for incorporation into the Wilderness System. The mineral surveys constitute one aspect of the suitability studies. The act directs that the results of such surveys are to be made available to the public and be submitted to the President and the Congress. This report discusses the results of a mineral survey of the Pond Mountain Roadless Area (8-035) and the Pond Mountain Addition Roadless Area (8-273), Cherokee National Forest, Carter County, Tennessee.

MINERAL RESOURCE POTENTIAL SUMMARY STATEMENT

The U.S. Geological Survey and U.S. Bureau of Mines studied the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee, to evaluate the mineral resource potential of the areas. The areas are on the northwestern flank of the Blue Ridge and are underlain by crystalline rocks of Precambrian age and clastic rocks and dolomite of Cambrian age.

A small amount of residual manganese and iron was mined near the northern boundary of the Pond Mountain Roadless Area. Iron prospects and thorium prospects in Precambrian rock were opened southeast of the area, but no ore has been mined from these deposits within the roadless area. The resource potential is low for manganese and iron in small areas in the northern part of the Pond Mountain Roadless Area.

The Precambrian rocks contain abundant tin, niobium, tungsten, beryllium, and a few other metals, which indicate that the rocks have been mineralized, most strongly in the Pond Mountain Addition Roadless Area. Most of the Pond Mountain Addition Roadless Area has low to high resource potential for tin, tungsten, and niobium. A small part of the Pond Mountain Roadless Area has low resource potential for tin, tungsten, and niobium. Beryllium, molybdenum, zinc, topaz, and fluorite might be recovered as byproducts of the mining of tin, tungsten, or niobium.

INTRODUCTION

The roadless areas encompass about 10 sq mi in the Cherokee National Forest in eastern Tennessee (fig. 1). They are in Carter County, about 7 mi southeast of Elizabethton and are near the Tennessee-North Carolina boundary. The area is heavily wooded and rises southward from an altitude of about 2,000 ft near Lake Watauga to about 4,000 ft in the Blue Ridge Mountains. Public roads surround the roadless areas, and they are crossed by the Appalachian Trail.

Geochemical fieldwork was done in 1980 by K. A. Duttweiler, W. R. Griffitts, and J. W. Whitlow of the U.S. Geological Survey. Prospects were examined and rocks and sediment sampled in 1981-83 by R. F. Bitar, P. T. Behum, and R. W. Hammack of the Bureau of Mines, who also mapped a shear zone that crosses Big Branch a short distance south of Pond Mountain. Supplementary fieldwork was done by M. L. Chatman. The broader features of the geology are from King and Ferguson (1960).

Little mining has been done in the roadless areas. A few prospects for radioactive materials were opened in the Precambrian rocks in and alongside the roadless areas in the 1950's, and iron prospects were opened in veins in the same general areas many years earlier. Manganese and iron were mined from residual deposits along the northern and northwestern edge of the areas.

The Federal government holds surface rights of about 99 percent of the roadless areas and mineral rights of about 54 percent. The rest of the rights are privately held. The U.S. Bureau of Land Management has issued oil and gas leases covering about 21 percent of the roadless areas or about 39 percent of the area for which the government holds mineral rights. In 1983, there were no active permits to prospect for solid minerals on government land.

MINING ACTIVITY

No mines or quarries were active in the vicinity of the roadless areas during the investigations in

NEW YORK
JUL 5 1989
STATE LIBRARY

best copy available



Figure 1.—Index map showing location of the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee.

1980–83. Prospecting and small-scale mining has been mainly of residual lateritic iron and manganese deposits and of veins that contain black iron oxides, uranium, and thorium (fig. 2).

GEOLOGY

The Pond Mountain area is underlain by a sequence of sedimentary rocks of early Paleozoic age that overlies Precambrian basement rocks.

The basement rocks are exposed in the southeastern part of the roadless areas and consist of Precambrian granitic rocks ranging in composition from granite to quartz monzonite to granodiorite, and smaller amounts of micaceous metamorphic rocks. Coarse, massive, pink granite is predominant, but small bands of finer grained red granite and fine-grained porphyritic granite are within the coarse granite, as are some steeply dipping phyllite or phyllonite layers.

Older basement rocks that have been thrust over the granitic rocks consist of a complex of quartz monzonite and granite that has been highly sheared, resulting in flaser gneiss, phyllonite, and mylonite. In addition, these sheared rocks were cut by pegmatites, quartz veins, and dikes. The metamorphism forming flaser gneiss, mylonite, and phyllite was plutonic and took place in Precambrian time. One shear zone in granitic rock was studied in detail. The thrusting, folding, and local shearing occurred after metamorphism in late Paleozoic time (King and Ferguson, 1960).

The sedimentary rocks lie unconformably on the basement rocks. They include, at the base, the thick Chilhowee Group of clastic rocks of Late Proterozoic to Early Cambrian age, overlain by a carbonate sequence, the Shady Dolomite.

The lowermost unit of the Chilhowee Group is the Unicoi Formation. It consists of thick-bedded quartzite, partly arkosic and conglomeratic, interbedded with some thin-bedded arkose.

The Hampton Formation, overlying the Unicoi Formation, is generally thought of as a shaly interval between the more sandy or quartzitic Unicoi and Erwin Formations below and above, respectively. It is composed mainly of dark-gray to greenish-gray shale and interbedded siltstone and arkosic sandstone in the lower part, and vitreous quartzite in the upper part.

The Erwin Formation is the uppermost unit of the Chilhowee Group and includes four members, which for this report have been grouped. They consist of interlayered white vitreous quartzite, dark ferruginous quartzite, siltstone, and shale. The white quartzite beds constitute a relatively small part of the formation but serve to distinguish it from the otherwise similar underlying Hampton Formation.

The Shady Dolomite of Early and Middle Cambrian age overlies the Chilhowee Group. It consists mostly of blue-gray and white dolomite but includes small amounts of limestone and a few beds of shale in the upper part. Dolomite of the Shady has been jointed, brecciated, and recrystallized. In many places, not shown on the map, the Shady Dolomite is covered by a waxy brown-buff residual clay formed by weathering of the dolomite. Concretions or lumps of manganese and iron oxides are in residuum overlying the lower part of the Shady Dolomite.

GEOCHEMISTRY

Concentrates of nonmagnetic heavy minerals panned from each stream in or near the roadless areas were analyzed mineralogically and spectrographically; most concentrates collected in the area of Precambrian rock contain concentrations of ore metals or minerals, and a few samples collected in the area of Paleozoic rocks contained ore metals or minerals (Duttweiler and others, 1987). Tin values exceed 1,000 ppm in all samples except one from the Precambrian rocks, niobium values are 200–2,000 ppm, tungsten 200–2,000 ppm, beryllium 10–200 ppm, and molybdenum 10–20 ppm. Most of these elements and fluorite and topaz are found in most of the world's leading tin-producing districts (Taylor, 1979).

The abundance of tin, beryllium, niobium, and tungsten associated with small amounts of molybdenum and zinc provide clear evidence that the Precambrian rocks in the study area have been mineralized. Topaz and fluorite in the heavy-mineral concentrates also indicate mineralization that is geochemically similar to that which formed tin and tungsten ores in many parts of the world (Taylor, 1979). Samples that are rich in niobium and tungsten are most abundant from the southeastern part of the study area, including samples containing topaz or fluorite. These higher values probably indicate that the mineralization was stronger in that general area. Columbite and wolframite, probably the main hosts of niobium and tungsten, respectively, do not survive transport over very long distances, so they may have disappeared from the sediment as it was reworked in the western and northern streams. More durable cassiterite, as well as phenakite, chrysoberyl, and some other beryllium minerals could persist for much longer distances, which would result in more widely dispersed tin and beryllium.

MINERAL DEPOSITS

Small residual manganese and iron deposits are known near the northern part of the roadless areas. The ore deposits consist of lumps of hydrous iron or manganese oxides embedded in tan to light-brown clay. These bodies are a residuum of weathering of the Shady Dolomite and are restricted to remnants of the Harrisburg or Valley Floor peneplain.

Black iron ore is exposed in the Precambrian rocks near, but not in, the roadless areas (Bayley, 1923). Possibly related uranium and thorium prospects were opened in, as well as near, the roadless areas. No minable deposits were found.

The investigations leading to this report have provided evidence of previously unknown mineralization with niobium, tin, tungsten, and beryllium. The veins of magnetite and black hematite in the Precambrian rocks and veins containing rare earths very likely were formed during the same mineralization. Although minor as iron resources, these veins are of use in indicating the extent of the mineralized area. These little known deposits of lithophile elements are speculative sources of tin, niobium, and tungsten. As such, they are more attractive exploration targets than the previously known residual iron and manganese deposits, which are known to be rather small. They are, accordingly, discussed in more detail.

best copy available



1. Unnamed Fe prospect
2. Carden Mn prospect
3. Grindstaff Mn prospect
4. Teaster and Ray Mn prospect
5. Unnamed Fe and Mn prospect
6. Cardens Bluff Fe prospect
7. Cardens Bluff Fe mine
8. Cardens Bluff Mn mine
9. Virginia Iron, Coal, and Coke Co. (C. A. Blevins) Fe, Mn prospects
10. Limonite Ore Bank, Fe
11. Unnamed Fe ore prospect
12. Unnamed Mn prospect
13. Sandstone quarry
14. Row Branch U prospect
15. Black Bear Fe prospect
16. Blacks Fe prospect
17. Rabbit Station Fe prospect
18. Mays Ridge Fe prospect
19. Lunsford Fe prospect
20. Finney and Teegarden Fe mine
21. Miller Fe prospect
22. Goodwin Field Branch U prospect
23. Laurel Cap U prospect
24. Wagener and McHone U prospect
25. Walnut Mountain U, Th, Nb, RE prospect
26. Unnamed Fe prospect
27. Granite quarry
28. Whitehead Fe prospect
29. Dennis Cove U prospect
30. Big Flats Branch U prospect
31. Sandstone quarry
32. Sand pit
33. Cedar Hill Mn mine

Figure 2.—Map showing mines and prospects in the Pond Mountain and Pond Mountain Addition Roadless Areas and vicinity. Mn, manganese; U, uranium; Nb, niobium; Ss, sandstone; Fe, iron; Th, thorium; RE, rare earths; Gr, granite.

Residual manganese and iron deposits

The roadless areas lie along the southeastern margin of the Hampton mining district. Minor amounts of brown iron ore and hard and soft manganese ores have been produced from residual clay of the Shady Dolomite. Most iron production took place during the middle to late 1800's and early 1900's. Many mines originally opened for iron were subsequently reopened for manganese production during World Wars I and II and during the Federal strategic stockpile purchase program between 1953 and 1959.

Individual manganese and iron deposits in the region are small and generally underlie less than 2 acres; most yielded no more than a few hundred tons of ore. The size and shape of the deposits in the residuum are irregular. Many are lenses that pinch out laterally but may be parts of series of lenses. Some areas that appear suitable for manganese and iron oxide accumulation contain no deposits. Although the Shady Dolomite is believed to be the source of the manganese and iron, no unusual concentrations of manganese have been found in unweathered dolomite (King and others, 1944).

Hard oxides, mainly psilomelane in nodules, are the most abundant manganese ore; soft oxides such as pyrolusite and wad are also present. The iron minerals consist mainly of limonite and goethite. Higher grade deposits of manganese oxides (about 40 percent Mn) and iron oxides (about 50 percent Fe) are more or less separate from one another. However, the lower grade deposits of either manganese or iron commonly contain several percent of the other metal. Manganese deposits in the Shady Dolomite have cobalt concentrations of as much as 1 percent (King and others, 1944), probably in lithiophorite. The highest cobalt value noted in a sample obtained near the study area is 0.10 percent in manganese ore from the Cardens Bluff mine (Bitar and others, 1985).

The parts of the study area with the largest deposits are along the lower slopes and base of the northwestern side of Pond Mountain where crests of low ridges representing the remnants of formerly extensive terraces are at two levels: between 2,080 and 2,200 ft in altitude, and between 2,260 and 2,380 ft in altitude. Smaller manganese deposits are on a series of smaller terrace remnants at a third level between 2,380 and 2,480 ft. Manganese nodules have washed downhill from the terraces to form small wash deposits in several places, mostly north of the roadless areas.

Some of the larger excavations for residual manganese or iron deposits are described below.

Cardens Bluff mine

The earliest iron mining near the areas was probably at the Cardens Bluff mine (loc. 7, fig. 2). Limonite and goethite were produced from an open cut and shafts in yellow residual clay.

The Cardens Bluff mine was reopened for manganese (loc. 8, fig. 2), and a large open cut yielded psilomelane and pyrolusite from residual clay. Manganese float was reported on a ridge crest southwest of the mine (King and others, 1944; loc. 5, fig. 2).

Exploration of the C. A. Blevins property on the ridge east of the Cardens Bluff mine in 1954 by Tennessee Manganese Corp., and in 1956 by Virginia Iron, Coal, and Coke Co., delineated a minable manganese deposit (U.S. Forest Service, written commun., 1981), but no mining was done.

Carden prospect

The Carden prospect (loc. 2, fig. 2) was explored at various times prior to 1941. Three cuts were made in residual clay containing hard manganese oxide nodules. In 1941, 3.6 long tons of manganese concentrates were produced (King and others, 1944).

Teaster and Ray prospect

The Teaster and Ray prospect (loc. 4, fig. 2) includes two open cuts and several prospect pits. First exploration was in 1918; in 1936, 5 long tons of manganese concentrates were produced (King and others, 1944; Bitar and others, 1985). Examination of the workings during the present study revealed pockets and lenses of hard manganese oxide nodules and of wad. Similar material was exposed in two older pits nearby.

Limonite Ore Bank

The Limonite Ore Bank (loc. 10, fig. 2) workings parallel the strike of the bedrock and consist of two trenches extending into open cuts. An unknown amount of iron ore was produced from residual clay between 1907 and 1909.

Mineral deposits in Precambrian rocks

Several types of mineral deposits have been explored in the Precambrian rocks, both in the southeastern part of the roadless areas and especially west and southeast of them. The commodities sought in these deposits were iron, uranium, and thorium. The present study disclosed, in the same area, unusual amounts of tin, niobium, beryllium, tungsten, fluorite, and topaz. Minerals containing niobium and rare earths were found in place in a shear zone. All of these metals and minerals may have been emplaced by the same episode of mineralization.

Magnetite-hematite prospects

Several iron-ore prospects in Precambrian granitic rocks south and southeast of the roadless areas (Bayley, 1923; Hamilton, 1940; fig. 2) were explored from about 1890 to 1912. The ore consisted of hematite, magnetite, or a mixture of both. No important iron deposits were found, but the prospecting showed black iron oxides to be a widespread product of mineralization.

The Whitehead prospect is on a ridge crest near the southern boundary of the Pond Mountain Addition Roadless Area (loc. 28, fig. 2). Bayley (1923) reported a thickness of 4 ft of flinty, blue-gray hematite, and granite containing a mixture of magnetite and hematite.

At the Miller prospect (loc. 21, fig. 2), surface and underground workings revealed crushed quartz syenite and crushed diorite containing magnetite, hornblende, and small veins of epidote and quartz. No

best copy available

commercial iron deposits were discovered, and the work was abandoned (Bayley, 1923).

Other magnetite and hematite occurrences include the Black Bear, Blacks, Rabbit Station, Mays Ridge, and Lunsford prospects; Finney and Teegarden mine; and an unnamed prospect (locs. 15-20 and 26, fig. 2).

Thorium and uranium deposits

Previous investigators reported only minor amounts of thorium and uranium minerals in veins and pegmatites in Precambrian granitic rocks near the roadless areas (locs. 14, 22-25, 29, and 30, fig. 2). Vein occurrences include reported "skarn" on Walnut Mountain that contains 0.09 percent U_3O_8 (Wagener and McHone, 1980) and anomalous concentrations of thorium and uranium at the Walnut Mountain, Goodwin Field Branch, and Laurel Gap prospects (S. W. Maher, Tennessee Division of Geology, oral commun., 1981). Occurrences in pegmatite include reported radiometric readings of four to ten times background values at the Dennis Cove prospect (Southern Interstate Nuclear Board, 1969), 0.021-0.15 percent U_3O_8 at the Row Branch prospect, and 0.057-0.22 percent U_3O_8 at the Big Flats Branch prospect (Stow, 1955).

A reconnaissance radiometric survey of the Precambrian crystalline rocks conducted during this study indicated that the total background radiation is 150-300 counts per second. Anomalies of two to three times background were detected in some granite and gneiss exposures. Higher radioactive anomalies of three to five times background were detected in the shear zone that contains ultramylonite and altered granitic rock.

Radiometric and fluorometric analyses of ultramylonite samples indicated an average of about 160 ppm thorium and 30 ppm uranium. The analyses revealed slightly anomalous thorium and uranium values in all granitic rock samples. A highly weathered phyllonite sample having radiation of three times background contained 79 ppm thorium and 13 ppm uranium.

Investigation of the Walnut Mountain prospect showed radiation of two to three times background. Analysis of medium-grained, veined granite indicated only 40 ppm thorium and 5 ppm uranium. Coarse-grained granite samples from the Laurel Gap and Goodwin Field Branch prospects contained 7 ppm thorium and 46 and 12 ppm uranium, respectively. Petrographic and X-ray microprobe examinations of samples of granite indicate that thorium and uranium are in monazite, thorite, and euxenite. Other accessory minerals include bastnaesite, xenotime, apatite, and zircon.

A little iron prospecting was done near a large fault zone south of Watauga Point (loc. 11, fig. 2). A pit 7-10 ft deep and 10-20 ft wide was excavated in sheared and fractured rock; veins developed along the fractures, one of which was 2-3 in. wide and contained uranium, thorium, niobium, and rare-earth elements (S. W. Maher, Tennessee Division of Geology, oral commun., 1982).

New Jersey Zinc Co., and H&H Co. opened several other prospects near the roadless areas. Among these, the Goodwin Field Branch and Laurel Gap prospects (locs. 22 and 23, fig. 2) were reported to be vein deposits (S. W. Maher, oral commun., 1982). Radioactive pegmatite outcrops were reported at the Dennis Cove prospect (Southern Interstate Nuclear Board, 1969) and at the Big Flats Branch and Row Branch prospects (Stow, 1955; locs. 14, 29, and 30, fig. 2).

Pechiney Ugine Kuhlmann Development, Inc., held uranium prospecting permits in the roadless areas from 1976 to 1979. Examination by this company of reported radioactive anomalies revealed no significant uranium deposits.

Deposits of niobium with rare metals

Shear zone deposits

During this investigation, Bitar and others (1985) mapped cataclastic rocks along a shear zone in the headwaters of Firescald and Big Branches that contain unusual amounts of niobium and rare earths. The shear zone, typified by distinctive outcrops of ultramylonite and by radiometric anomalies in residual soils, trends northwest, about normal to thrust faults bordering regional allochthons. Observed dips range from 45° to 60° to the northeast.

Ultramylonite, the core of the shear zone, is bounded by alternating layers of granite, altered granite, and phyllonite. X-ray fluorescence analysis of 15 ultramylonite samples indicated an average niobium content of about 0.1 percent. Niobium is rather uniformly distributed along the length of the ultramylonite body. Petrographic and X-ray microprobe analyses revealed that niobium occurs in columbite, fergusonite, euxenite, and pyrochlore in a matrix of finely brecciated quartz and iron-rich muscovite. Other minerals include zircon, thalenite, thorium- and nonthorium-bearing monazite, thorite, bastnaesite, and cerianite. Hematite, in fine-grained clusters of grains and in fracture fillings, is ubiquitous and imparts a faint, dark-red tinge to fresh ultramylonite surfaces. Limonite pseudomorphs after pyrite occur in clusters and thin bands.

Altered granite and phyllonite of the outer part of the shear zone have niobium contents that range from 0.005 to 0.104 percent. Chemical and petrographic analyses of green altered granite revealed that niobium-bearing and other accessory minerals are similar to those in the adjacent ultramylonite but in smaller amounts.

Vein deposits

Two rare-metal occurrences were found near Walnut Mountain by regional uranium and thorium prospecting. Both occurrences are veins along fractures or foliation.

A vein at locality 24 (fig. 2) was described by Wagener and McHone (1980) as an altered zone of foliated granite containing medium-grained skarn. They reported that the material contained 0.2 percent niobium, 0.2 percent yttrium, 0.1 percent zirconium, and 0.02 percent lanthanum. No skarn was found here during our study. However, 0.08 percent gadolinium, 0.04 percent tin, 6 ppm tungsten, and 26 ppm beryllium were detected in a sample of foliated granite.

Niobium and rare-earth elements were also detected in a 2-3-in.-wide vein at the Walnut Mountain prospect (loc. 25, fig. 2) (S. W. Maher, Tennessee Division of Geology, oral commun., 1982).



EXPLANATION OF RESOURCE POTENTIAL

The presence of three or more commodities in the group tin, tungsten, niobium, fluorite, and topaz indicates high resource potential; the presence of two indicates moderate potential, and the presence of one indicates low potential.

HIGH RESOURCE POTENTIAL

Tin, tungsten, and niobium

MODERATE RESOURCE POTENTIAL

Tin, tungsten, and niobium

LOW RESOURCE POTENTIAL

Tin, tungsten, and niobium

Manganese and iron

Figure 3.--Map showing mineral resource potential in the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee.

best copy available

Placer deposits

Placer deposits of niobium, tin, zirconium, rare-earths, thorium, and uranium minerals may exist in the roadless areas. High concentrations of niobium, tungsten, tin, beryllium, and rare earths, and lower concentrations of thorium and uranium, were detected in panned concentrates. The rare earths are in monazite, bastnaesite, xenotime, and euxenite.

Sand and stone

Stone for local road construction was provided by the sandstone quarry in the Unicoi Formation, a small granite quarry, and a small sandstone quarry in the Erwin Formation (locs. 13, 27, and 31, fig. 2). None of these quarries have been operated in recent years. A sand pit, also inactive (loc. 32, fig. 2), is in weathered rock of the Hesse Quartzite. Operations were terminated because, at shallow depth, the rock became too hard to crush for sand (S. W. Maher, oral commun., 1982).

ASSESSMENT OF MINERAL RESOURCE POTENTIAL

Resources in crystalline rocks

Within the roadless areas, the Precambrian rocks were mineralized, and the intensity of mineralization in a particular area is indicated by the number of ore metals and minerals in the group consisting of tin, tungsten, niobium, fluorite, and topaz. The presence of three or more of these commodities indicates high resource potential, the presence of two indicates moderate potential, and the presence of one indicates low potential (fig. 3). Zinc, molybdenum, or beryllium commonly accompany these metals and minerals (Duttweiler and others, 1987) but were not used in classifying the mineral resource potential of any areas. Most of the Pond Mountain Addition Roadless Area has low to high resource potential for tin, tungsten, and niobium. A small part of the Pond Mountain Roadless Area has low resource potential for tin, tungsten, and niobium.

A mineralized shear zone that crosses Big Branch and Firescald Branch near their heads was studied in detail. Having an average thickness of 6 ft and a length of 3,750 ft, it contains, above altitude 3,040 ft, about 17.5 million cubic feet, or 1.5 million short tons of rock. The estimated amounts of various metals in this body are given in table 1.

Resources in residual material

Areas having potential resources of manganese and iron along the northern boundary of the Pond Mountain Roadless Area are (1) remnants of prominent terraces that are underlain by Shady Dolomite and (2) hillslopes mantled with colluvium below the terraces. Estimates of resources are based only on lumps of hard manganese oxides in the clay, even though small masses of soft pyrolusite and wad in the deposits were sought during early mining because of their high manganese content. The resource potential is low for manganese and iron in small areas in the northern part of the Pond Mountain Roadless Area.

Other resources

Clay, sand, and stone are present in the roadless areas. Clay might be a byproduct of the mining of residual manganese ore. Sand deposits are too small for practical exploration, and stone is overlain by extensive weathered overburden and has no advantage over deposits that are closer to major markets.

Table 1.--Niobium and other rare metals in a mineralized shear zone in Pond Mountain Addition Roadless Area, Tennessee

Metal	Grade (percent)[1]	Amount (in millions of lbs of contained metal)[1]
Niobium.....	0.1	3.1
Tantalum.....	.01	0.31
Zirconium.....	.36	11
Yttrium.....	.06	1.8
Ytterbium.....	.03	0.92
Tin.....	.05	1.5
Thorium.....	.016	0.49
Uranium.....	.003	0.09

[1]Bitar and others (1985, p. 39, 54).

REFERENCES

Bayley, W. S., 1923, The magnetic iron ores of east Tennessee and western North Carolina: Tennessee Division of Geology Bulletin 29, 252 p.

Bitar, R. F., Behum, P. T., Hammack, R. W., and Chatman, M. L., 1985, Minerals investigation of Pond Mountain and Pond Mountain Addition RARE II Further Planning Areas, Carter County, Tennessee: U.S. Bureau of Mines Open-File Report MLA 8-85, 54 p.

Duttweiler, K. A., Griffitts, W. R., Cooley, E. F., and Whitlow, J. W., 1987, Geochemical map of the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee: U.S. Geological Survey Miscellaneous Field Studies Map MF-1648-B, scale 1:24,000.

Hamilton, S. H., 1940, Report on the iron ore bodies of Carter County, Tennessee, and Avery County, North Carolina, contiguous to eastern Tennessee and Western North Carolina Railroad: unpublished file material for War Minerals Reports, available at U.S. Bureau of Mines, Intermountain Field Operations Center, Denver, CO, 68 p.

King, P. B., and Ferguson, H. W., 1960 [1961], Geology of northeasternmost Tennessee, with a section on Description of the basement rocks, by Warren Hamilton: U.S. Geological Survey Professional Paper 311, 136 p.

King, P. B., Ferguson, H. W., Craig, L. C., and Rodgers, John, 1944, Geology and manganese deposits of northeastern Tennessee: Tennessee Division of Geology Bulletin 52, 275 p.

Southern Interstate Nuclear Board, 1969, Uranium in the southern United States: U.S. Atomic Energy Commission WASH-1128, 230 p.

Stow, H. M., 1955, Report of the radiometric reconnaissance in Virginia, North Carolina, eastern Tennessee, and parts of South Carolina, Georgia, and Alabama: U.S. Atomic Energy Commission report RME-3107, 33 p.

Taylor, R. G., 1979, Geology of tin deposits: Amsterdam, Elsevier Scientific Publishing Co., 543 p.

Wagener, H. D., and McHone, J. G., 1980, Uranium mineralization in the Wilson Creek and Cranberry gneisses and the Grandfather Mountain Formation, North Carolina and Tennessee: South Portland, Maine, Chiasma Consultants, Inc., 25 p.

best copy available

DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

Note: The following correlation and description are for the geologic base map printed in gray

CORRELATION OF MAP UNITS

Qal	Holocene	QUATERNARY
€r	Middle and Lower Cambrian	CAMBRIAN
€s		
€e	Lower Cambrian	
€Zh	Lower Cambrian or Late Proterozoic	CAMBRIAN AND PROTEROZOIC
Zu		LATE PROTEROZOIC
pEg		PRECAMBRIAN
pEc / pEu		

DESCRIPTION OF MAP UNITS

Qal ALLUVIUM
€r ROME FORMATION (MIDDLE AND LOWER CAMBRIAN)--Maroon or brown shale interbedded with shaly dolomite
€s SHADY DOLOMITE (MIDDLE AND LOWER CAMBRIAN)--Blue-gray and white dolomite
€e ERWIN FORMATION (LOWER CAMBRIAN)--White to buff quartzite, siltstone, and shale
€Zh HAMPTON FORMATION (LOWER CAMBRIAN OR LATE PROTEROZOIC)--Dark-gray to greenish-gray shale, minor quartzite
Zu UNICOI FORMATION (LATE PROTEROZOIC)--Pink to brown quartzite, partly arkosic or conglomeratic, minor dark-green shale
pEg GRANITE, QUARTZ MONZONITE, AND GRANODIORITE (PRECAMBRIAN)--Granite is pink and massive
pEc COARSE GRANITE AND QUARTZ MONZONITE (PRECAMBRIAN)--Much sheared to form flaser gneiss and phyllonite
pEu METAMORPHIC AND IGNEOUS ROCKS, UNDIVIDED (PRECAMBRIAN)

CONTACT--Dashed where approximately located; dotted where concealed
FAULT--Showing dip where known and relative horizontal movement. Dashed where approximately located; dotted where concealed. D, downthrown side; U, upthrown side
THRUST FAULT--Showing dip where known. Sawteeth on upper plate; dashed where approximately located; dotted where concealed

STRIKE AND DIP OF BEDS
Inclined
Overturned
Vertical
Horizontal
STRIKE AND DIP OF FOLIATION
STRIKE AND DIP OF CLEAVAGE
Inclined
Vertical
APPROXIMATE BOUNDARY OF ROADLESS AREAS

IRON MOUNTAIN FAULT
Pond Mountain Roadless Area
Pond Mountain Addition Roadless Area
Mineralized shear zone Mapped by Bitar and others (1985)

82°10′ 82°5′ 82°00′
36°22′30″ 36°20′ 36°15′

Base from U.S. Geological Survey Elizabethton, Watauga Dam, 1960

Geology modified from King and Ferguson (1960)

SCALE 1:24 000
1 MILE
1 KILOMETER

CONTOUR INTERVAL 40 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

MINERAL RESOURCE POTENTIAL MAP OF THE POND MOUNTAIN AND POND MOUNTAIN ADDITION ROADLESS AREAS, CARTER COUNTY, TENNESSEE

By
W.R. Griffitts, M.L. Chatman, and R.F. Bitar

1989

MISCELLANEOUS FIELD STUDIES
MAP MF-1648-A
PAMPHLET ACCOMPANIES MAP



Geology modified from King and Ferguson (1960)

EAS, CARTER COUNTY, TENNESSEE

EXPLANATION OF RESOURCE POTENTIAL

The presence of tin, tungsten, niobium, fluorite, and topaz was used as the basis for determining the low to high resource potential for tin, tungsten, and niobium. The presence of three or more of these metals or minerals indicates high resource potential; the presence of two indicates moderate potential; and the presence of one indicates low potential.

HIGH RESOURCE POTENTIAL

Tin, tungsten, and niobium

MODERATE RESOURCE POTENTIAL

Tin, tungsten, and niobium

LOW RESOURCE POTENTIAL

Tin, tungsten, and niobium

Manganese and iron

STUDIES RELATED TO WILDERNESS

The Wilderness Act (Public Law 88-577, September 3, 1964) and related acts require the U.S. Geological Survey and the U.S. Bureau of Mines to survey certain areas on Federal lands to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a mineral survey of the Pond Mountain Roadless Area (8-035) and the Pond Mountain Addition Roadless Area (8-273), Cherokee National Forest, Carter County, Tennessee. The Pond Mountain and Pond Mountain Addition Roadless Areas were classified as further planning areas during the Second Roadless Area Review and Evaluation (RARE II) by the U.S. Forest Service, January 1979.

MINERAL RESOURCE POTENTIAL SUMMARY STATEMENT

The U.S. Geological Survey and U.S. Bureau of Mines studied the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee, to evaluate their mineral resource potential. The roadless areas are on the northwestern flank of the Blue Ridge and are underlain by crystalline rocks of Precambrian age and clastic rocks and dolomite of Cambrian age.

A small amount of residual manganese and iron was mined near the northern boundary of the roadless areas. Iron prospects and thorium prospects were opened southeast of the areas, but no ore has been mined within the roadless areas themselves. The resource potential is low for manganese and iron in small areas in the northern part of the Pond Mountain Roadless Area.

The Precambrian rocks contain abundant tin, niobium, tungsten, beryllium, and a few other metals, which indicate that the rocks have been mineralized, most strongly in the Pond Mountain Addition Roadless Area. Most of the Pond Mountain Addition Roadless Area has low to high resource potential for tin, tungsten, and niobium. A small part of the Pond Mountain Roadless Area has low resource potential for tin, tungsten, and niobium. Beryllium, molybdenum, zinc, topaz, and fluorite might be recovered as byproducts of the mining of tin, tungsten, or niobium.

INTRODUCTION

The roadless areas encompass about 10 sq mi in the Cherokee National Forest in eastern Tennessee (fig. 1). They are in Carter County, about 7 mi southeast of Elizabethton, and are near the Tennessee-North Carolina boundary. The area is heavily wooded and rises southward from an altitude of about 2,000 ft near Lake Watauga to about 4,000 ft in the Blue Ridge Mountains.

Geochemical fieldwork was done in 1980 by K. A. Duttweiler, W. R. Griffitts, and J. W. Whitlow of the U.S. Geological Survey. Examination of prospects and sampling of rocks and stream sediment was done in 1981-1983 by R. F. Bitar, P. T. Behum, and R. W. Hammack of the U.S. Bureau of Mines; they also mapped a shear zone that crosses Big Branch a short distance south of Pond Mountain.

Little mining has been done in the roadless areas. A few prospects for radioactive materials were opened in the Precambrian rocks in and alongside the Pond Mountain Addition Roadless Area in the 1950's, and iron prospects were opened in veins in the same general areas many years earlier. Manganese and iron were mined from residual deposits along the northern and northwestern part of the Pond Mountain Roadless Area. Information about the broader features of geology is from a report by King and Ferguson (1960).

GEOLOGY

The Pond Mountain area is underlain by a sequence of sedimentary rocks of early Paleozoic age that overlie Precambrian basement rocks. The basement rocks in the southeastern part of the roadless areas consist of Precambrian granitic rocks and smaller amounts of micaceous metamorphic rocks. Coarse, massive, pink granite is predominant, but small layers of finer grained red granite and fine-grained porphyritic granite are within the coarse granite as are some steeply dipping phyllite or phyllonite layers.

Older basement rocks that have been thrust over the granitic rocks are quartz monzonite and granite that have been highly sheared, resulting in laser gneiss, phyllonite, and mylonite (map unit pCc) cut by pegmatites, quartz veins, and dikes. One shear zone in granitic rock was studied in detail. The sedimentary rocks that lie unconformably on the basement rocks are the thick Chilhowee Group of clastic rocks of Late Proterozoic to Early Cambrian age, and the overlying Shady Dolomite (unit Cs) of Early and Middle Cambrian age.

The lowest unit of the Chilhowee Group is the Unicoi Formation (unit Cu) of Late Proterozoic age. It consists of thick-bedded quartzite, partly arkosic and conglomeratic, interbedded with minor thin-bedded arkose.

The Hampton Formation (unit Ch) of Late Proterozoic or Early Cambrian age is a shaly interval between the more sandy or quartzitic Unicoi and Erwin Formations below and above, respectively. It is mainly dark-gray to greenish-gray shale and interbedded siltstone and arkosic sandstone in the lower part and vitreous quartzite in the upper part.

The Erwin Formation (unit Ce) of Early Cambrian age is the top unit of the Chilhowee Group and includes interlayered, white, vitreous quartzite, dark, ferruginous quartzite, siltstone, and shale. The white quartzite beds are a small part of the formation and distinguish it from the otherwise-similar, underlying Hampton Formation.

The Shady Dolomite, of Early and Middle Cambrian age, consists mostly of blue-gray and white dolomite but includes a little limestone and a few beds of shale in the upper part. The dolomite has been jointed, brecciated, and recrystallized. In many places, the Shady Dolomite is covered by a waxy, brown-buff, residual clay. Concretions or lumps of manganese and iron oxides are in this residuum.

GEOCHEMISTRY

Heavy-mineral concentrates panned from streams in or near the roadless areas were analyzed mineralogically and spectrographically. Most concentrates collected in the area of Precambrian rock contained concentrations of ore metals or minerals, but only a few samples collected in the area of Paleozoic rocks did (Duttweiler and others, 1987). Tin values exceed 1,000 ppm in all samples except one from the Precambrian rocks, niobium values are 200-2,000 ppm, tungsten, 200-2,000 ppm, beryllium, 10-200 ppm, and molybdenum, 10-20 ppm. Most of these elements and fluorite and topaz are found in most of the world's leading tin-producing districts (Taylor, 1979).

The abundance of tin, beryllium, niobium, and tungsten associated with small amounts of molybdenum, zinc, topaz, and fluorite provide clear evidence that the Precambrian rocks in the roadless areas have been mineralized. Sediment samples rich in niobium and tungsten are most abundant from the southeastern part of the Pond Mountain Addition Roadless Area, indicating that the mineralization was stronger there.

MINERAL DEPOSITS

Deposits were opened in the nineteenth century in search of black iron ore in the Precambrian rocks southeast of the roadless areas, and several prospects were opened in the 1950's for uranium and thorium, also in the Precambrian rocks in and near the roadless areas. No mineable deposits were found.

The investigations leading to this report have provided evidence of previously unknown mineralization that introduced niobium, tin, tungsten, fluorine, and beryllium. The veins of magnetite and black hematite in the Precambrian rocks very likely were formed during the same mineralization and therefore may be of use in indicating the extent of the mineralized areas. These little-known deposits of lithophile elements are speculative sources of industrial raw materials and are more attractive exploration targets than the previously known small iron and manganese deposits.



Figure 1.--Index map showing location of the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee.

Residual iron and manganese deposits

The roadless areas lie along the southeastern margin of the Hampton mining district, from which minor amounts of brown iron ore and hard and soft manganese ores have been produced from residual clay of the Shady Dolomite. Most iron mining took place during the middle to late 1800's and early 1900's. Many mines originally opened for iron were subsequently reopened for manganese production during World Wars I and II and between 1953 and 1959.

Individual manganese and iron deposits generally underlie areas of less than 2 acres; most yielded no more than a few hundred tons of ore. Many are lenses that pinch out laterally but may be parts of series of lenses.

Hard oxides, mainly psilomelane in nodules, are the most abundant manganese ore, soft oxides such as pyrolusite and wad are also present. The iron minerals are mainly limonite and goethite. Higher grade ores of manganese oxides, containing about 40 percent Mn, or of iron oxides, containing about 50 percent Fe, are more or less separate from one another, whereas the lower grade ores of either manganese or iron commonly contain several percent of the other metal. Manganese deposits in the Shady Dolomite have cobalt concentrations of as much as 1 percent (King and others, 1944), probably in lithiophorite. The highest cobalt content noted in a sample obtained near the study area is 0.10 percent in manganese ore from the Cardens Bluff mine (Bitar and others, 1985).

The largest deposits are along the lower parts and base of the northwestern side of Pond Mountain where the crests of low ridges, representing the remnants of formerly extensive terraces, are at two levels: between 2,080 and 2,200 ft in altitude, and between 2,260 and 2,380 ft in altitude. Smaller manganese deposits are on a series of smaller terrace remnants at a third level between 2,380 and 2,480 ft. Manganese nodules have washed downhill from the terraces to form small colluvial deposits in several places, mostly north of the roadless areas.

Some of the larger excavations for residual manganese or iron are described below.

Cardens Bluff mine

The earliest iron mining near the study area was probably at the Cardens Bluff mine (loc. 7, fig. 2). Limonite and goethite were produced from an open cut and shafts in yellow residual clay.

The Cardens Bluff mine was reopened for manganese (loc. 8, fig. 2), and a large open cut yielded psilomelane and pyrolusite from residual clay. Manganese oxide float was reported on a ridge crest southwest of the mine (King and others, 1944; loc. 5, fig. 2). Exploration of the ridge east of the Cardens Bluff mine in 1954 by Tennessee Manganese Corp., and in 1956 by Virginia Iron, Coal, and Coke Co., delineated a minable manganese deposit (U.S. Forest Service, written commun., 1981), but no mining was done.

Carden prospect

The Carden prospect (loc. 2, fig. 2) was explored at various times prior to 1941 and, in 1941, 3.6 long tons of manganese concentrates were produced (King and others, 1944).

Toaster and Ray prospect

The Toaster and Ray prospect (loc. 4, fig. 2) includes two open cuts and several prospect pits. In 1938, 5 long tons of manganese concentrates were produced (King and others, 1944; Bitar and others, 1985). Examination of the workings during the present study revealed pockets and lenses of hard manganese oxide nodules and of wad. Similar material was exposed in two older pits nearby.

Limonite Ore Bank

The Limonite Ore Bank (loc. 10, fig. 2) workings parallel the strike of the bedrock and consist of two trenches extending into open cuts. An unknown amount of iron ore was produced from residual clay between 1907 and 1909.

Mineral deposits in Precambrian rocks

Several types of mineral deposits have been explored for iron, uranium, and thorium in the Precambrian rocks mostly in the Pond Mountain Addition Roadless Area and especially south and east of the roadless areas. In the same area the present study disclosed unusual amounts of tin, niobium, beryllium, tungsten, fluorite, and topaz. Minerals containing niobium and rare earths were found in place in a shear zone.

Magnetite-hematite prospects

Several iron-ore prospects in Precambrian granitic rocks south and southeast of the roadless areas (Bayley, 1923; Hamilton, 1940, fig. 2) were explored from about 1890 to 1912. The ore consisted of hematite, magnetite, or a mixture of both. No important iron deposits were found, but the prospecting showed black iron oxides to be a widespread product of mineralization.

The Whitehead prospect is on a ridge crest just outside the southern boundary of the Pond Mountain Addition Roadless Area (loc. 28, fig. 2). Bayley (1923) reported a thickness of 4 ft of flinty, blue-gray hematite and granite containing a mixture of magnetite and hematite.

At the Miller prospect (loc. 21, fig. 2), surface and underground workings revealed crushed quartz syenite and crushed diorite containing magnetite, hornblende, and small veins of epidote and quartz. No commercial iron deposits were discovered, and the work was abandoned (Bayley, 1923).

Other magnetite and hematite occurrences include the Black Bear, Blacks, Rabbit Station, Mays Ridge, and Lunsford prospects, the Finney and Teagarden mine, and an unnamed prospect (loc. 15-20 and 26, fig. 2).

Thorium and uranium deposits

Previous investigators reported only minor amounts of thorium and uranium in veins and pegmatites in Precambrian granitic rocks near the roadless areas (locs. 14, 22-25, 29, and 30, fig. 2). Vein occurrences include "skarn" on Walnut Mountain that contains 0.09 percent U_3O_8 (Wagener and McHone, 1980) and rather low concentrations of thorium and uranium at the Walnut Mountain, Goodwin Field Branch, and Laurel Gap prospects (S. W. Maher, Tennessee Division of Geology, oral commun., 1981). Occurrences in pegmatite include reported radiometric readings of 4-10 times background values at the Dennis Cove prospect (Southern Interstate Nuclear Board, 1969), 0.021-0.15 percent U_3O_8 at the Row Branch prospect, and 0.057-0.22 percent U_3O_8 at the Big Flats Branch prospect (Stow, 1955).

A reconnaissance radiometric survey of the Precambrian crystalline rocks conducted during this study indicated that the total background radiation is 150-300 counts per second. Anomalies of two to three times background were detected in some granite and gneiss exposures. In the Pond Mountain Addition Roadless Area, higher radioactive anomalies of three to five times background were detected in the shear zone that contains ultramylonite and altered granitic rock.

Radiometric and fluorometric analyses of ultramylonite samples indicated an average of about 160 ppm thorium and 30 ppm uranium. The analyses revealed slightly anomalous thorium and uranium values in all granitic rock samples. A highly weathered phyllonite sample having radiation of three times background contained 79 ppm thorium and 13 ppm uranium.

Investigation of the Walnut Mountain prospect showed radiation of two to three times background. Analysis of medium-grained, veined granite indicated only 40 ppm thorium and 5 ppm uranium. Coarse-grained granite samples from the Laurel Gap and Goodwin Field Branch prospects contained 7 ppm thorium and 46 and 12 ppm uranium, respectively.

Petrographic and X-ray microprobe examinations of samples of granite from the roadless areas indicate that thorium and uranium are in monazite, thorite, and euxenite. Other accessory minerals include bastnaesite, xenotime, apatite, and zircon.

A little iron prospecting was done near a large fault zone south of Watauga Point (loc. 11, fig. 2). A pit 7-10 ft deep and 10-20 ft wide was excavated in sheared and fractured rock. It exposed veins developed along the fractures, one of which was 2-3 in. wide and contained uranium, thorium, niobium, and rare-earth elements (S. W. Maher, Tennessee Division of Geology, oral commun., 1982).

New Jersey Zinc Co. and H&H Co. opened several other prospects near the roadless areas. Among these, The Goodwin Field Branch and Laurel Gap prospects (locs. 22 and 23, fig. 2) were reported to be vein deposits (S. W. Maher, oral commun., 1982). Radioactive pegmatite outcrops were reported at the Dennis Cove prospect (Southern Interstate Nuclear Board, 1969) and at the Big Flats Branch and Row Branch prospects (Stow, 1955, locs. 14, 29, and 30, fig. 2).

Pechiney Ugine Kuhlman Development, Inc., held uranium prospecting permits in the roadless areas from 1976 to 1979 but found no significant uranium deposits.

Deposits of rare metals

Shear-zone deposits

During this investigation, Bitar and others (1985) mapped cataclastic rocks along a shear zone in the headwaters of Firescald and Big Branches that contain unusual amounts of niobium and rare earths. The shear zone trends northwest, about normal to thrust faults that border the regional allochthons. Observed dips in the shear zone range from 45° to 60° northeast.

Ultramylonite, the core of the shear zone, is bounded by alternating layers of granite, altered granite, and phyllonite. X-ray fluorescence analyses of 15 ultramylonite samples indicated an average niobium content of about 0.1 percent and that niobium is rather uniformly distributed along the length of the ultramylonite body. Petrographic and electron microprobe analyses revealed that niobium occurs in columbite, fergusonite, euxenite, and pyrochlore, in a matrix of finely brecciated quartz and iron-rich muscovite. Other minerals include zircon, thalenite, thorium- and nonthorium-bearing monazite, thorite, bastnaesite, and cerianite. Hematite is ubiquitous and imparts a faint, dark-red tinge to fresh ultramylonite surfaces. Limonite pseudomorphs after pyrite are in clusters and thin bands.

Altered granite and phyllonite of the outer part of the shear zone have niobium contents that range from 0.005 to 0.104 percent. Niobium-bearing and other accessory minerals in green, altered granite are similar to those in the adjacent ultramylonite, but they are in smaller amounts.

Vein deposits

Rare metals were found at two places near Walnut Mountain during regional uranium and thorium prospecting. At both places veins are along fractures or foliation.

A vein at locality 24 (fig. 2) was described by Wagener and McHone (1980) as an altered zone of foliated granite containing medium-grained skarn. They reported that the material contained 0.2 percent niobium, 0.2 percent yttrium, 0.1 percent zirconium, and 0.02 percent lanthanum. No skarn was found here during our study. However, 0.08 percent gadolinium, 0.04 percent tin, 6 ppm tungsten, and 26 ppm beryllium were detected in a sample of foliated granite.

Niobium and rare-earth elements were also detected in a 2-3-in.-wide vein at the Walnut Mountain prospect (loc. 25, fig. 2) (S. W. Maher, Tennessee Division of Geology, oral commun., 1982).

Placer deposits

Placer deposits of niobium, tin, zirconium, rare-earth minerals, thorium, and uranium may exist in the roadless areas, but possible placer deposits are small because of the small size of most of the gravel beds.

Sand and stone

A sand pit, two sandstone quarries, and a granite quarry near the roadless areas have not been operated in recent years. The sand pit (loc. 32, fig. 2) is in weathered rock of the Hesse Quartzite. Operations were terminated because, at shallow depth, the rock became too hard to process as sand (S. W. Maher, oral commun., 1982). The sandstone quarry in the Unicoi Formation, a small granite quarry, and a small sandstone quarry in the Erwin Formation (locs. 13, 27, and 31, fig. 2) provided stone for local use.

ASSESSMENT OF MINERAL RESOURCE POTENTIAL

Resources in crystalline rocks

Within the roadless areas, the crystalline Precambrian rocks were mineralized, and the intensity of mineralization is indicated by the number of metals and minerals found in the group consisting of tin, tungsten, niobium, fluorite, and topaz. The presence of one or more of these metals or minerals was used as the basis for determining the low to high resource potential for tin, tungsten, and niobium. The presence of three of these commodities indicates high resource potential, the presence of two indicates moderate potential, and the presence of one indicates low potential. Zinc, molybdenum, or beryllium commonly accompany these metals and minerals (Duttweiler and others, 1987) but were not used in classifying the mineral resource potential of any areas. Most of the Pond Mountain Addition Roadless Area has low to high resource potential for tin, tungsten, and niobium. A small part of the Pond Mountain Roadless Area has low resource potential for tin, tungsten, and niobium.

A mineralized shear zone that crosses Big Branch and Firescald Branch near their heads was studied in detail. Having an average thickness of 6 ft and a length of 3,750 ft, it contains, above an altitude of 3,040 ft, about 17.5 million cubic feet, or 1.5 million short tons of rock. Its estimated content of various metals is given in table 1.

Table 1.--Niobium and other rare metals in a mineralized shear zone in Pond Mountain Addition Roadless Area

(Shear zone shown on map)

Metal	Grade (percent)[1]	Amount (millions of pounds of contained metal)
Niobium	0.1	3.1
Tantalum	.01	0.31
Zirconium	.36	11
Yttrium	.06	1.8
Ytterbium	.03	0.92
Tin	.05	1.5
Thorium	0.016	.49
Uranium	.003	.09

[1]Bitar and others (1985, p. 39, 54).

Resources in residual material

Areas having potential resources of manganese and iron along the northern boundary of the Pond Mountain Roadless Area are (1) remnants of prominent terraces that are underlain by Shady Dolomite and (2) hillslopes mantled with manganiferous colluvium below the terraces. Estimates of resource potential are based on lumps of hard manganese oxides in the clay, even though small masses of soft pyrolusite and wad in the deposits were sought during early mining because of their high manganese content. The resource potential is low for manganese and iron in small areas in the northern part of the Pond Mountain Roadless Area.

Other resources

Clay, sand, and stone are present in the roadless areas. Clay might be a byproduct of the mining of residual manganese ore. Sand deposits are too small for practical exploitation, and stone is overlain by extensive weathered overburden and has no advantage over deposits that are closer to major markets.

REFERENCES

Bayley, W. S., 1923, The magnetic iron ores of east Tennessee and western North Carolina: Tennessee Division of Geology Bulletin 29, 254 p.

Bitar, R. F., Behum, P. T., Hammack, R. W., and Chatman, M. L., 1985, Mineral investigation of Pond Mountain and Pond Mountain Addition RARE II Further Planning Areas, Carter County, Tennessee: U.S. Bureau of Mines Open-File Report MLA 6-85, 54 p.

Duttweiler, K. A., Griffitts, W. R., Cooley, E. F., and Whitlow, J. W., 1987, Geochemical map of the Pond Mountain and Pond Mountain Addition Roadless Areas, Carter County, Tennessee: U.S. Geological Survey Miscellaneous Field Studies Map MF-1648-B, scale 1:24,000.

Hamilton, S. B., 1940, Report on the iron ore bodies of Carter County, Tennessee, and Avery County, North Carolina, contiguous to eastern Tennessee and Western North Carolina Railroad: unpublished file material for War Minerals Reports, available at U.S. Bureau of Mines, Intermountain Field Operations Center, Denver, CO, 68 p.

King, P. B., and Ferguson, H. W., 1960 [1961], Geology of northeasternmost Tennessee, with a section on Description of the basement rocks, by Warren Hamilton: U.S. Geological Survey Professional Paper 311, 136 p.

King, P. B., Ferguson, H. W., Craig, L. C., and Rodgers, John, 1944, Geology and manganese deposits of northeastern Tennessee: Tennessee Division of Geology Bulletin 52, 275 p.

Southern Interstate Nuclear Board, 1969, Uranium in the southern United States: U.S. Atomic Energy Commission WASH-1128, 230 p.

Stow, H. M., 1955, Report of the radiometric reconnaissance in Virginia, North Carolina, eastern Tennessee, and parts of South Carolina, Georgia, and Alabama: U.S. Atomic Energy Commission report RME-3107, 33 p.

Taylor, R. G., 1979, Geology of tin deposits: Amsterdam, Elsevier Scientific Publishing Co., 543 p.

Wagener, H. D., and McHone, J. G., 1980, Uranium mineralization in the Wilson Creek and Cranberry gneisses and the Grandfather Mountain Formation, North Carolina and Tennessee: South Portland, Maine, Chlaems Consultants, Inc., 25 p.



1. Unnamed Fe prospect
2. Carden Mn prospect
3. Grindstaff Mn prospect
4. Toaster and Ray Mn prospect
5. Unnamed Fe and Mn prospect
6. Cardens Bluff Fe prospect
7. Cardens Bluff Fe mine
8. Cardens Bluff Mn mine
9. Virginia Iron, Coal, and Coke Co. (Co. A, Blevins) Fe, Mn prospects
10. Limonite Ore Bank, Fe
11. Unnamed Fe ore prospect
12. Unnamed Mn prospect
13. Sandstone quarry
14. Row Branch U prospect
15. Black Bear Fe prospect
16. Blacks Fe prospect
17. Rabbit Station Fe prospect
18. Mays Ridge Fe prospect
19. Lunsford Fe prospect
20. Finney and Teagarden Fe mine
21. Miller Fe prospect
22. Goodwin Field Branch U prospect
23. Laurel Gap U prospect
24. Wagener and McHone U prospect
25. Walnut Mountain U, Th, Nb, RE prospect
26. Unnamed Fe prospect
27. Granite quarry
28. Whitehead Fe prospect
29. Dennis Cove U prospect
30. Big Flats Branch U prospect
31. Sandstone quarry
32. Sand pit
33. Cedar Hill Mn mine

Figure 2.--Map showing mines and prospects in and near the Pond Mountain and Pond Mountain Addition Roadless Areas. Fe, iron; Gr, granite; Mn, manganese; Nb, niobium; RE, rare earths; Ss, sandstone; Th, thorium; U, uranium.

Manuscript approved for publication December 20, 1988

PRINTED 1989

For sale by U.S. Geological Survey, Map Distribution, Box 25286, Federal Center, Denver, CO 80225

best copy available